

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>C07H 21/00, C12N 15/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 91/14699</b> <b>(43) International Publication Date:</b> 3 October 1991 (03.10.91)
<b>(21) International Application Number:</b> PCT/US91/01893 <b>(22) International Filing Date:</b> 21 March 1991 (21.03.91)  <b>(30) Priority data:</b> 496,852                      21 March 1990 (21.03.90)                      US  <b>(71) Applicant:</b> THE UNIVERSITY OF COLORADO FOUNDATION, INC. [US/US]; P.O. Box 1140, Boulder, CO 80306 (US).  <b>(72) Inventors:</b> CECH, Thomas, R. ; 1545 Rockmont Circle, Boulder, CO 80303 (US). HERSCHLAC, Daniel ; 3135 Arnett Avenue, Apt. 301, Boulder, CO 80304 (US).  <b>(74) Agent:</b> WARBURG, Richard, J.; 611 West Sixth Street, 34th Floor, Los Angeles, CA 90017 (US).		<b>(81) Designated States:</b> AT (European patent), AU, BB, BE (European patent), BG, BR, CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC, MG, MW, NL (European patent), NO, PL, RO, SD, SE (European patent), SU.  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> SITE SPECIFIC CLEAVAGE OF SINGLE-STRANDED DNA  <b>(57) Abstract</b>  Method for specifically cleaving a single stranded DNA molecule. The method includes providing an RNA molecule having a deoxyribonuclease activity independent of any protein, and contacting that RNA molecule with the single stranded DNA molecule to cause the single stranded DNA molecule to be cleaved.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

- 1 -

## SITE SPECIFIC CLEAVAGE OF SINGLE-STRANDED DNA

Background of the Invention

This invention was made with support from a grant  
5 from the National Institute of Health (Grant No. GM  
28039). The U.S. Government has rights in the invention.

This invention relates to methods for cleaving  
single stranded deoxyribonucleic acid (DNA).

Zaug and Cech, 231 Science 470, 1986, state that  
10 certain ribonucleic acid (RNA) molecules, now called  
ribozymes, are able to catalyze cleavage and rejoining of  
single stranded RNA molecules. This activity appears to  
be specific for ribonucleotides; no reaction is observed  
with deoxyribonucleotides. A chain of five  
15 deoxyribocytosine residues (dC<sub>5</sub>), however, is a  
competitive inhibitor of cleavage of a chain of five  
cytosine residues (C<sub>5</sub>).

Cech, 236 Science 1532, 1987, describes cleavage  
of mixed oligonucleotides (containing both ribo- and  
20 deoxyribo-nucleotides), but only cleavage between a  
deoxyribonucleotide and a ribonucleotide is observed. No  
cleavage is observed between two deoxyribonucleotide  
residues.

Sugimoto et al, 17 Nucleic Acids Research, 355,  
25 368, 1989, describe a reaction where a circular RNA  
molecule is opened by addition of a dinucleotide, which  
may include one or more deoxynucleotides. The  
dinucleotide is added to the RNA molecule, and the circle  
opened to form a single fragment. No cleavage of DNA is  
30 described.

Summary of the Invention

The invention features a method for specifically  
cleaving a single stranded DNA molecule. The method  
includes providing an RNA molecule (or ribozyme) having a  
35 deoxyribonuclease activity independent of any protein,

- 2 -

and contacting that RNA molecule with the single stranded DNA molecule to cause the single stranded DNA molecule to be cleaved. In this method, a linear DNA molecule is cleaved to form two or more DNA fragments, and a circular  
5 DNA molecule is cleaved to form one or more DNA fragments. These fragments are released from the ribozyme as free molecules.

By "specifically cleaving" is meant that the RNA molecule is active at one or only a few short nucleotide sequences of a single stranded DNA molecule. Such  
10 sequences have a length between three and eight deoxynucleotides, generally about four to six deoxynucleotides. The specific sequence that is cleaved can be chosen by altering a portion of the active site of  
15 the RNA molecule, a well known procedure described, for example, by Cech et al., PCT application WO88/04300; Haseloff and Gerlach, 334 Nature 585, 1988; Been and Cech, 47 Cell 207, 1986; and Haseloff et al., PCT Application WO89/05852. A single stranded DNA molecule  
20 includes any molecule having a contiguous sequence of at least four, preferably eight, deoxynucleotides. Such molecules may be bonded with other DNA or RNA molecules, or hybrids thereof.

In preferred embodiments, the method includes  
25 providing the RNA molecule in a reaction medium at a concentration sufficient to cause cleavage of at least 1% of a population of the DNA molecules in an hour, most preferably at a concentration sufficient to cause cleavage of at least 10% of such DNA molecules. The  
30 reaction medium generally includes DNA molecules at any desired concentration, e.g., between 0.1nM and 10 $\mu$ M, and is maintained at a pH between 6.0 and 9.5, and at a temperature between 20°C and 70°C. The RNA molecule is generally provided at a concentration between 0.1 $\mu$ M and  
35 100 $\mu$ M.

- 3 -

The RNA molecule is preferably one that cleaves an RNA molecule to leave a 3'-hydroxyl group; most preferably the RNA molecule includes the deoxyribonuclease cleaving active site of an RNA molecule chosen from a group I or group II intron (Lambowitz, 56 Cell 323, 1989; Peebles et al., 44 Cell 213, 1986; Van der Veen et al., 44 Cell 225, 1986; and Jarrell et al., 263 J. Biol. Chem. 3432, 1988), or RNase P (Altman, Advances in Enzymology and related areas of Molecular Biology, 1, 1989), or derivatives thereof. Even more preferably, the RNA molecule is a derivative of an intervening sequence (IVS) of Tetrahymena, e.g., T. thermophila, and the RNA molecule is L-19, L-21, or a derivative thereof.

By "derivative" is meant any RNA molecule which has the active site of any known ribozyme which has a deoxyribonuclease activity. This active site may be altered to specifically cleave a desired single-stranded DNA sequence. Such an RNA molecule need only contain those essential portions of the ribozyme necessary for the deoxyribonuclease activity. Such ribozymes can be readily designed by those of ordinary skill in the art by use of any number of standard techniques.

In other preferred embodiments, the contacting step further includes providing magnesium ions, providing guanosine, guanosine monophosphate, guanosine diphosphate or guanosine triphosphate or another guanosine containing moiety, e.g., a dinucleotide containing guanosine; and treating the DNA molecule with a denaturant, e.g., glyoxylate or glyoxal, prior to the contacting step; and the method further includes providing a chelator of magnesium ions, e.g., EDTA, to stop further reaction of the RNA molecule with the DNA molecule.

- 4 -

In yet other preferred embodiments, the RNA molecule includes a binding site for single stranded DNA which is complementary to a cleavage site on the single stranded DNA molecule; the method includes providing a plurality of RNA molecules, each having a different binding site for single-stranded DNA; and the method further includes separating the cleaved DNA fragments by electrophoresis within a gel matrix.

This invention provides a method by which single stranded DNA molecules can be specifically cleaved. Such a method is useful because it allows the mapping of specific cleavage sites or nucleotide sequences within DNA. Prior methods for determination of specific DNA sites included digestion of double-stranded DNA with restriction endonucleases. Such restriction endonucleases are active at specific sequences, but their specificity cannot be readily altered. In addition, there are many DNA sequences which no restriction endonuclease is known to cleave. In contrast, ribozymes can be readily altered or synthesized to cleave any desired single-stranded DNA sequence, and thus can be used to characterize DNA molecules with only few limitations on the sites that are recognized.

Although the method is limited to cleavage of single-stranded DNA, double-stranded DNA can be used in the method if it is first denatured to form single stranded DNA. This method is extremely useful in vitro for genome mapping of humans or other organisms. In addition, the ribozymes can be used in vivo to cleave single stranded DNA. For example, ribozymes having high activity for cleavage of single stranded DNA can be used to cleave single stranded DNA viruses, and thus reduce their infectivity and any disease symptom that they cause.

- 5 -

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

#### Description of the Preferred Embodiments

5 The drawings will first briefly be described.

#### Drawings

Fig. 1 is a diagrammatic representation of a proposed deoxyribonuclease reaction of a ribozyme;

10 Figs. 2 and 3 are photographs of polyacrylamide gels demonstrating the deoxyribonuclease activity of a ribozyme (in Fig. 2 the lanes are referred to below as lanes 1-8 starting from the left hand side); and

Fig. 4 is a graphical representation of the time course for reaction of a ribozyme with a single-stranded  
15 DNA substrate.

#### Ribozymes

As discussed above, any ribozyme having deoxyribonuclease activity may be used in this invention. Generally, useful ribozymes are those that  
20 cleave an RNA molecule to leave a 3'-hydroxyl group, or a 3' phosphate, and not a 2', 3' cyclic phosphate. These ribozymes can be isolated, and modified to provide derivatives, as described by Cech et al., PCT, supra; Lambowitz, supra, and Van der Veen, supra. Any of a  
25 broad group of derivatives of naturally-occurring ribozymes are useful in this invention, all of which can be readily isolated by those of ordinary skill in the art. For example, as described by Cech et al., PCT, supra, the ribozyme of Tetrahymena thermophilla can be  
30 readily modified so that its active site is active to cleave RNA molecules of different RNA sequence. In a similar manner, this ribozyme can be altered to cleave DNA molecules having differing DNA sequence. Such alterations can be performed by site-directed  
35 mutagenesis, or equivalent procedure (see, .g., Murphy

- 6 -

and Cech, Proc. Nat. Acad. Sci. USA., 1990). Generally, the active site is constructed to have a base sequence complementary to the base sequence to be cleaved; that is, at least 3 of 4 bases are positioned to form base pairs, e.g., A is complementary to, and will base pair with, T or U, and G is complementary to, and will base pair with, C or U (where A, T, U, C and G are the standard symbols for adenine, thymine, uracil, cytosine and guanine respectively). In addition, deletion or addition of various portions of the ribozyme, or mutation by other means, can be used to create smaller or larger ribozymes which retain, and preferably increase, their deoxyribonuclease activity. Such alterations are within the ordinary skill of one in the art, and no undue experimentation is required to determine which of those ribozymes are active. Such ribozymes need merely be created and tested in a DNA cleavage protocol, for example, as described below, to determine whether or not they have sufficient deoxyribonuclease activity to be useful in this invention.

Generally, ribozymes useful in this invention are those which can be provided at a concentration in a reaction medium sufficient to cause at least 1%, preferably 10% or more, of a population of DNA molecules to be cleaved within an hour, under the conditions described below. The ratio of the concentration of ribozyme to the concentration of DNA can be varied dependent upon the ribozyme and the DNA and can be in any desired ratio, generally it is between  $1:10^6$  and 1000:1, preferably between 1:10 and 1000:1. The higher the ratio of ribozyme to DNA the faster the reaction will proceed. Higher ratios, however, are wasteful of the ribozyme and need not be used. It is important that the ribozyme used be designed such that the single stranded DNA substrate is cleaved into one, two or more fragments, which are



- 7 -

released from the ribozyme. It is also important that the ribozyme not be altered by the process so that it can recycle and cleave additional DNA molecules.

For in vivo use, it is desirable to have ribozymes that have higher deoxyribonuclease activity under the conditions prevailing in vivo. To this end, ribozymes having an activity to cleave at least 10% of a population of DNA molecules within an hour at 37°C are desirable.

Although alteration of a synthetic version of an RNA occurring naturally in Tetrahymena thermophila is described above, this example is not limiting in the invention since those of ordinary skill in the art can readily alter other group I or group II ribozymes, or RNase P to provide ribozymes useful in this invention.

There follow examples of the use of ribozymes in vitro for cleavage of single stranded DNA. These examples are not limiting to the invention, are provided merely to illustrate the invention, and those skilled in the art will readily recognize that the invention can be practiced as broadly claimed below.

Referring to Fig. 1, applicants propose a general model for the cleavage of single stranded DNA by a ribozyme. This model is meant only to illustrate applicant's present understanding of the reaction, and is not meant to be limiting to the invention. This example relates specifically to the deoxyribonuclease activity of a synthetic RNA version of a RNA in Tetrahymena thermophila ribozyme; this synthetic version is a ribozyme. In Fig. 1, the deoxyribonuclease reaction of the L-21 ScaI ribozyme (Zaug et al., 27 Bioc. 8924, 1988) with a DNA substrate 18 (deoxy-CCCUCUA<sub>5</sub>) is shown. All of the steps are reversible. The binding of guanosine (G) prior to oligonucleotide substrate 18 is shown for simplicity; the two binding sites (substrate and G binding sites) are essentially independent so that either

- 8 -

order of addition can occur. The 5' sequence of the ribozyme (shaded, GGGAGG-5'), called the 5' exon-binding site 12, is responsible for recognition of the substrate by base pairing.

5 Referring again to Fig. 1, it is proposed that ribozyme 10 has a substrate binding site 12, and a guanosine binding site 14. A guanosine residue (G) binds to site 14. Active site 12 of the ribozyme then interacts with a substrate 18 to form the structure shown  
10 as 20. Ribozyme 10 in structure 20 causes cleavage of substrate 18 to release a deoxy A<sub>5</sub> chain having the guanosine residue attached (GdA<sub>5</sub>), and the remainder of the substrate (dCCCUCU<sub>OH</sub>), to reform ribozyme 10. Thus, ribozyme 10 acts in a catalytic manner since it is able  
15 to act on any number of substrates to cause cleavage of each substrate and subsequent release of the fragments resulting from the cleavage.

L-21 ScaI of T. thermophila

Referring to Fig. 2, there is shown the results of  
20 the deoxynucleotide cleavage reaction of a DNA oligonucleotide substrate catalyzed by the ribozyme L-21 ScaI. The ribozyme was formed by standard procedure (Zaug et al., 27 Bioc. 8924, 1988). The substrate, deoxyCCCUCUA<sub>5</sub>, was provided at a concentration of  
25 approximately 2 nanomolar and labeled at its 5'-end with <sup>32</sup>P. It was synthesized on an Applied Biosciences DNA synthesizer. This substrate was incubated at a concentration of 2 nanomolar with 1 micromolar L-21 ScaI ribozyme (id.) for 15 hours at 50°C in the presence of  
30 800 micromolar G, 50 millimolar MES (buffer), pH 7.0, and 10 millimolar MgCl<sub>2</sub> (lane 4). The same reaction was performed without G (lane 5), with 800 micromolar ATP replacing the G (lane 6), without MgCl<sub>2</sub> (lane 7), and without ribozyme (lane 8). Lane 2 shows the substrate  
35 alone, and lane 3 shows the putative product, deoxy <sup>32</sup>P-

- 9 -

CCCUCU. Lane 1 shows the products from digestion of the substrate by 0.2 units per microliter P1 nuclease in 1 millimolar Tris, pH 7.5, 2 millimolar zinc chloride, 0.1 millimolar EDTA, and 0.1 microgram per microliter tRNA for 60 minutes at 37°C.

Ribozyme reactions were initiated by addition of the DNA substrate after a 10 minute preincubation of the ribozyme, buffer, and  $MgCl_2$  at 50°C. Reactions were quenched with EDTA and urea at 0°C and subjected to electrophoresis on the 20% denaturing polyacrylamide gel, as described by Zaug et al., id.

The DNA substrate was specifically cleaved at the same position as the analogous RNA substrate for this ribozyme. A single labelled product is formed from this DNA substrate. This product comigrates with deoxy  $^{32}P$ -CCCUCU (lane 3) and with the appropriate product from P1 nuclease digestion of the DNA substrate (lane 1).  $Mg^{2+}$  is required (lane 7). ATP (adenosine triphosphate) cannot substitute for G (lane 6), and GTP (guanosine triphosphate) can substitute for G (not shown), each of which is also a property of the reaction with RNA substrates. A small amount of product is formed in the absence of added G (lanes 5 and 6, seen in longer exposures). This is analogous to the slow hydrolysis of RNA substrates that occurs in the absence of G. Similar experiments with deoxy- $^{32}P$ -CCCUCUA also gave specific cleavage between U and A dependent on the presence of the ribozyme.

Referring to Fig. 3, using DNA substrate that was 3'-end-labeled, it was demonstrated that the guanosine cofactor was covalently added to the 3'-product fragment. Deoxy-CCCUCUA<sub>5</sub>  $^{32}P$ -3'dA (2nM, produced by reaction of deoxy-CCCUCUA<sub>5</sub> with [ $\alpha$ - $^{32}P$ ]3'dATP in the presence of terminal transferase) was incubated for four hours with 4  $\mu M$  L-21 ScaI ribozyme, 50 mM MES, pH 7.0,

- 10 -

and 10 or 100 mM  $\text{MgCl}_2$ . One aliquot was quenched with EDTA and urea (shown as "-" in Fig. 3), the other was treated with 0.05 U/ $\mu\text{l}$  ribonuclease T1 for 1 hour at 37°C (shown as "+" in Fig. 3). Lanes 3 and 4 show reaction with 1 mM G and 10 mM  $\text{MgCl}_2$ ; lanes 5 and 6 show reaction with 1  $\mu\text{M}$  G and 100 mM  $\text{MgCl}_2$ ; lanes 11 and 12 show reaction with 1mM GTP and 100 mM  $\text{MgCl}_2$  (reaction is faster with higher  $\text{MgCl}_2$  concentrations); lanes 7 and 8 are from incubation with 1 mM GTP and 100 mM  $\text{MgCl}_2$  in the absence of ribozyme; lanes 1 and 10 show substrate alone; and lanes 2 and 9 show deoxy- $\text{A}_5$   $^{32}\text{P}$ -3'dA(M). The product from reaction of deoxy-CCCUCUA $_5$   $^{32}\text{P}$ -3'dA with GTP (lane 11) migrated faster in a polyacrylamide gel than the product from reaction with G (lanes 3 and 5, conditions as above; no products were observed without a guanosine cofactor, without  $\text{MgCl}_2$ , or without ribozyme). The products from the ribozyme reactions with GTP and with G (lanes 4, 6 and 12) comigrated following treatment by ribonuclease T1, which cleaves after (ribo)-guanosine residues that have 3'-phosphoryl groups. The new product formed from T1 treatment comigrated with  $\text{HOdA}_5$   $^{32}\text{P}$ 3'dA $_{\text{OH}}$  (lanes 2 and 9), as expected from the mechanism outlined in Figure 1.

The DNA endonuclease reaction is catalytic. For example, a reaction mixture with 25  $\mu\text{M}$  DNA substrate (deoxy-CCCUCUA $_5$ ) and a trace (about 1nM) of 5'-end-labeled substrate was incubated with 0.55  $\mu\text{M}$  ribozyme and 800  $\mu\text{M}$  G for 20 hours. Five percent of the labeled DNA was converted to product, which corresponds to more than two turnovers.

Referring to Fig. 4, there is shown a time course for formation of product (deoxy  $^{32}\text{P}$ -CCCUCU) from deoxy  $^{32}\text{P}$ -CCCUCUA $_5$  (2 nM) with 1.1 micromolar ribozyme, 800 micromolar G, 50 millimolar MES, pH 7, and 10 millimolar

- 11 -

MgCl<sub>2</sub> at 50°C. The reactions were quenched, products separated by denaturing gel electrophoresis, and quantitated with an AMBIS radioanalytical scanner. As can be seen from the graph, approximately 1% of product is released every 30-50 minutes under these conditions.

DNA substrates have a lower affinity and a slower rate of reaction than a corresponding RNA substrate with a ribozyme. Lengthening of the internal guide sequence will improve binding, and may increase the rate of DNA endonuclease reaction.

#### Uses

As discussed above, the method of this invention is useful for determining the location of specific DNA sequences in a single stranded DNA molecule. Such analysis is analogous to restriction endonuclease mapping, using restriction endonuclease proteins and double stranded DNA. Generally, double stranded DNA is denatured, and the single stranded DNA molecules resulting from the denaturation, contacted with a ribozyme under appropriate conditions (e.g., those described above) to produce one, two or more single stranded DNA fragments which can then be separated in a gel matrix. The size of the resulting fragments reflects the position of the specific DNA site in the single stranded DNA molecule.

Separate aliquots of the DNA may be treated with different ribozymes so that the location of different cleavage sites can be determined. In some experiments it may be useful to contact the DNA with two, or even more, ribozymes; or the DNA may be contacted with one ribozyme and after removing the first ribozyme, contacted with a second ribozyme.

The method of this invention can be used in vivo by contacting single stranded DNA, within a cell or in medium surrounding the cell, with a ribozyme that is

- 12 -

active under in vivo conditions. Alternatively, the  
ribozyme may be synthesized in vivo by transfection of a  
DNA molecule into a cell in a manner that causes  
expression of that DNA molecule to produce the desired  
5 ribozyme in vivo. That ribozyme may then be active to  
cleave any single stranded DNA within the cell.

Other embodiments are within the following claims.

- 13 -

Claims

- 1           1. A method for specifically cleaving a single  
2 stranded DNA molecule, comprising the steps of:  
3           (a) providing an RNA molecule having a  
4 deoxyribonuclease activity independent of any protein,  
5 and  
6           (b) contacting said RNA molecule with said single  
7 stranded DNA molecule to cause said single stranded DNA  
8 molecule to be cleaved.
- 1           2. The method of claim 1, further comprising  
2 providing said RNA molecule in a reaction medium at a  
3 concentration sufficient to cause cleavage of at least 1%  
4 of a population of the DNA molecules in an hour.
- 1           3. The method of claim 1, further comprising  
2 providing said RNA molecule in a reaction medium at a  
3 concentration sufficient to cause cleavage of at least  
4 10% of a population of the DNA molecules in an hour.
- 1           4. The method of claim 1, further comprising  
2 selecting said RNA molecule from one that cleaves an RNA  
3 molecule to leave a 3'-hydroxyl group.
- 1           5. The method of claim 4, wherein said DNA  
2 molecule comprises the deoxyribonuclease cleaving active  
3 site of an RNA molecule chosen from a group I or group II  
4 intron, or RNase P, or derivatives thereof.
- 1           6. The method of claim 4, wherein said RNA  
2 molecule is a derivative of an intervening sequence of  
3 Tetrahymena.

- 14 -

1           7. The method of claim 5, wherein said RNA  
2 molecule is L-19, L-21, or a derivative thereof.

1           8. The method of claim 1, wherein said contacting  
2 step further comprises providing  $Mg^{2+}$  ions.

1           9. The method of claims 1, wherein said  
2 contacting step further comprises providing guanosine or  
3 guanosine triphosphate.

1           10. The method of claim 8, further comprising the  
2 step of providing a chelator of said  $Mg^{2+}$  ions to stop  
3 further reaction of said RNA molecule with said DNA  
4 molecule.


1           11. The method of claim 10, wherein said chelator  
2 is ethylenediaminetetraacetic acid (EDTA).

1           12. The method of claim 1, wherein said RNA  
2 molecule comprises a binding site for single stranded  
3 DNA, which binding site is complementary to nucleotides  
4 adjacent to a cleavage site on said single stranded DNA  
5 molecule.

1           13. The method of claim 12, further comprising  
2 providing a plurality of RNA molecules having different  
3 binding sites for single stranded DNA.

1           14. The method of claim 12, further comprising  
2 separating the two or more DNA fragments by  
3 electrophoresis within a gel.





154

2/3




FIG.2




FIG.3




FIG.4

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/US91/01893

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC(5): C07H 21/00; C12N 15/00		
U.S. CL.: 586/27, 28, 29; 435/91, 173, 199		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
U.S.	536/27, 28, 29; 435/6, 91, 173, 199	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched *		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT <sup>8</sup></b>		
Category *	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
X,P	Biotechnology, Volume 8, issued 09 April 1990, John Hodgson, "New Ways To Cleave Nucleic Acids", pages 348 and 349, see entire document.	1-14
X,P	Nature, Volume 344, issued 29 March 1990, Robertson et al., "Selection <u>in vitro</u> of an RNA enzyme that specifically cleaves single-stranded DNA". pages 467 and 468, see entire document.	1-14
X,P	Nature, Volume 345, issued 14 June 1990, Geoffrey North, "Expanding the RNA repertoire", pages 576-578, see entire document.	1-14
<p>* Special categories of cited documents: <sup>10</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
12 July 1991	26 JUL 1991	
International Searching Authority	Signature of Authorized Officer	
ISA./US	James O. Wilson	